

Chiral Zero Modes on Intersecting Heterotic 5-branes

Tetsuji Kimura* and Shun'ya Mizoguchi†

*Theory Center, High Energy
Accelerator Research Organization (KEK)
Tsukuba, Ibaraki 305-0801, Japan*

(Dated: Dec.8, 2009)

Abstract

We show that there exist two **27** and one $\overline{\mathbf{27}}$ of E_6 , net one chiral supermultiplet as zero modes localized on the intersecting 5-branes in the $E_8 \times E_8$ heterotic string theory. A heterotic background is constructed by the standard embedding in the smeared solution, and the Dirac equation is solved explicitly on this background. It provides, after a compactification of some of the transverse dimensions, a Randall-Sundrum II like brane-world set-up in heterotic string theory.

PACS numbers: 11.25.Mj, 11.25.Wx, 11.30.Qc

*Electronic address: tetsuji@post.kek.jp

†Also at Department of Particle and Nuclear Physics, The Graduate University for Advanced Studies.;

Electronic address: mizoguch@post.kek.jp

In this letter, we construct an intersecting 5-brane solution in the $E_8 \times E_8$ heterotic string theory by the so-called standard embedding in the known smeared intersecting NS5-brane solution of type II theories. We then study the zero modes of the relevant Dirac operator on this background and show that there exist three localized chiral zero modes, two of which are in the **27** representation of E_6 , and one in the $\overline{\mathbf{27}}$ representation. They give rise to net *one* **27** of massless chiral fermions in the four-dimensional spacetime. Therefore, this is the first example of a brane set-up in heterotic string theory that supports four-dimensional chiral matter fermions transforming as an E_6 gauge multiplet [11].

This letter is a concise version of the paper [1], to which the reader is referred for further detail.

We start with the neutral smeared solution [2]:

$$\begin{aligned}
ds^2 &= \sum_{i=0,7,8,9} \eta_{ij} dx^i dx^j + h(x^1)^2 \sum_{\mu=1,2} \delta_{\mu\nu} dx^\mu dx^\nu + h(x^1) \sum_{\mu=3,4,5,6} \delta_{\mu\nu} dx^\mu dx^\nu, \\
e^{2\phi} &= h(x^1)^2, \\
H_{\mu\nu\lambda} &= \begin{cases} \frac{h'}{2} \left(= \frac{\xi |x^1|'}{2} \right) & \text{if } (\mu, \nu, \lambda) = (2, 3, 4), (2, 5, 6) \text{ or their even permutation,} \\ -\frac{h'}{2} \left(= -\frac{\xi |x^1|'}{2} \right) & \text{if } (\mu, \nu, \lambda) = (2, 4, 3), (2, 6, 5) \text{ or their even permutation,} \\ 0 & \text{otherwise,} \end{cases} \quad (1)
\end{aligned}$$

where

$$h(x^1) = h_0 + \xi |x^1|. \quad (2)$$

All other components of H_{MNL} vanish. h_0 and ξ are real constants. The prime ' denotes the differentiation with respect to x^1 , and $|x^1|'$ is therefore a step function. This is a solution to equations of motion of the leading-order NSNS-sector Lagrangian in type II theories. The solution describes a pair of intersecting NS5-branes [3, 4, 5] stretching in dimensions as shown in Table I. These branes are delocalized in the x^2, x^3, x^4, x^5 and x^6 directions. Consequently, the solution depends only on x^1 , and hence the name ‘‘smeared solution’’.

TABLE I: Dimensions in which the 5-branes stretch.

	0	1	2	3	4	5	6	7	8	9
5-brane1	○					○	○	○	○	○
5-brane2	○			○	○			○	○	○




FIG. 1: $h(x)$ with $\xi < 0$. (a) The brane has a positive tension. The string coupling decreases linearly from a positive value h_0 , to necessarily cross the x^1 axis, where the string coupling becomes zero. We identify this point as the “end of the world”. (b) By a change of the coordinate the points $x = \pm \frac{h_0}{|\xi|}$ are mapped to $z = \pm\infty$. The profile of $h(x(z))$ becomes similar to the warp factor of the RS II model.

The coefficient ξ in the definition $h(x^1)$ is related to the tension of the brane. It turns out that the brane tension V is related to ξ as [1]

$$\xi = -\kappa^2 V h_0^{\frac{5}{2}}, \quad (3)$$

where κ is the Newton constant.

If $\xi = 0$, the solution is reduced to a flat Minkowski space. If $\xi > 0$, the brane tension is negative. It is doubtful whether such an object may consistently exist in heterotic string theory. It is also puzzling that the string coupling becomes stronger as one goes away from the branes. Thus, we assume that $\xi < 0$, then the brane has a positive tension. The string coupling $h(x^1)$ is now convex upwards in x^1 , and decreases linearly from a positive value h_0 , to necessarily cross the x^1 axis, where the string coupling becomes zero. Beyond that point, $h(x^1)$ becomes negative (FIG.1(a)). We identify this point as the “end of the world”; one can send this point infinitely far away by the coordinate transformation

$$z = -\text{sign}(x^1) \log \frac{h(x^1)}{h_0}, \quad (4)$$

where z is the new coordinate (FIG.1(b)). Then the function $h(x^1)$ is expressed simply as

$$h = h_0 e^{-|z|}. \quad (5)$$

Apparently, this looks similar to the Randall-Sundrum (RS) II model [6], but there are the following differences: The first is that we have no bulk cosmological constant. Secondly,

as we see in a moment, there exist chiral zero modes on the branes, which are in the **27** representation of E_6 . This is not an assumption but a logical consequence of string theory. The final difference is in the warp factor. Unlike the RS models, our four-dimensional metric is not warped in the string frame [12].

We now construct an intersecting solution in the $E_8 \times E_8$ heterotic string theory by the standard embedding, similarly to the symmetric 5-brane [5]. The (generalized) spin connections [7, 8, 9] of the neutral intersecting background are identified as the gauge connection $A_\mu^{\alpha\beta} = (\omega + H)_\mu^{\alpha\beta}$. The result is

$$\begin{aligned}
A_{\mu=1}^{\alpha\beta} &= 0, \\
A_{\mu=2}^{\alpha\beta} &= \frac{h'}{h} \begin{pmatrix} -s & & \\ & \frac{1}{2}s & \\ & & \frac{1}{2}s \end{pmatrix} = \frac{h'}{h} \left(-\frac{3\lambda_3 + \sqrt{3}\lambda_8}{4} \right) \otimes s, \\
A_{\mu=3}^{\alpha\beta} &= \frac{h'}{2h^{\frac{3}{2}}} \begin{pmatrix} & -\mathbf{1} & \\ & & \\ \mathbf{1} & & \end{pmatrix} = \frac{h'}{2h^{\frac{3}{2}}} (-i\lambda_2) \otimes \mathbf{1}, \\
A_{\mu=4}^{\alpha\beta} &= \frac{h'}{2h^{\frac{3}{2}}} \begin{pmatrix} & -s & \\ -s & & \\ & & \end{pmatrix} = \frac{h'}{2h^{\frac{3}{2}}} (-\lambda_1) \otimes s, \\
A_{\mu=5}^{\alpha\beta} &= \frac{h'}{2h^{\frac{3}{2}}} \begin{pmatrix} & -\mathbf{1} & \\ & & \\ & & \mathbf{1} \end{pmatrix} = \frac{h'}{2h^{\frac{3}{2}}} (-i\lambda_5) \otimes \mathbf{1}, \\
A_{\mu=6}^{\alpha\beta} &= \frac{h'}{2h^{\frac{3}{2}}} \begin{pmatrix} & -s & \\ & & \\ -s & & \end{pmatrix} = \frac{h'}{2h^{\frac{3}{2}}} (-\lambda_4) \otimes s,
\end{aligned} \tag{6}$$

where λ_i 's ($i = 1, \dots, 8$) are the Gell-Mann matrices and $\mathbf{1} \equiv \begin{pmatrix} 1 & \\ & 1 \end{pmatrix}$, $s \equiv i\sigma_2 = \begin{pmatrix} & 1 \\ -1 & \end{pmatrix}$.

The explicit expressions of $\omega_\pm$ show [1] that both are $SU(3)$ connections. Since we have embedded ω_+ into the gauge connection A , the Bianchi identity is reduced to $dH = 0$, and




FIG. 2: Broken generators which give rise to zero modes.

the solution (1) is consistent with it. An $SU(3)$ piece of the $E_8(\times E_8)$ gauge connection is given a nonzero expectation value. On the other hand, the fact that $\omega_- \in SU(3)$ implies that the Killing spinor equations for the gravitino as well as gaugino variations have a common single Killing spinor. It can be checked that this also satisfies the equation for the dilatino SUSY variation to lowest order. Thus, the background (1) together with (6) preserve 1/4 of supersymmetries.

In the case of a single symmetric 5-brane [5], the connection ω_+ embedded was in $SU(2)$, and the unbroken gauge symmetry was the centralizer E_7 . In the present intersecting case, the connection embedded into E_8 is in $SU(3)$, and therefore the unbroken gauge symmetry is E_6 . The adjoint representation of E_8 is decomposed into

$$\mathbf{248} = (\mathbf{78}, \mathbf{1}) \oplus (\mathbf{27}, \mathbf{3}) \oplus (\overline{\mathbf{27}}, \overline{\mathbf{3}}) \oplus (\mathbf{1}, \mathbf{8}) \quad (7)$$

as representations of the subalgebra $E_6 \times SU(3)$. Since the $E_8 \times E_8$ gauge field A_M has by construction a vev in $SU(3)$, the latter three gauge rotations are the moduli (FIG.2).

Let us focus on the E_6 non-singlet moduli. The spontaneously broken generators in $(\mathbf{27}, \mathbf{3}) \oplus (\overline{\mathbf{27}}, \overline{\mathbf{3}})$ give rise to Nambu-Goldstone bosons, each of which has one bosonic degree of freedom. On the other hand, since a $D = 4$, $\mathcal{N} = 1$ chiral supermultiplet needs *two* bosonic degrees of freedom, the Nambu-Goldstone bosons which transform as $\mathbf{27}$ and $\overline{\mathbf{27}}$ must be combined to form a single $\mathcal{N} = 1$ chiral supermultiplet. That is, the E_6 non-singlet moduli form *three* chiral supermultiplets in the $\mathbf{27}$ (or $\overline{\mathbf{27}}$, but not both) representation of E_6 .

At first sight, one might think that the argument above would be contradictory to the

well-known fact in Calabi-Yau compactifications that the number of chiral generations are determined by the Dirac index, in which the same decomposition (7) is used and *one triplet* of zero modes together corresponds to *one* supermultiplet, and is not counted as three. Of course, it is not a contradiction, because what we consider here is not the fermionic zero modes of the Dirac operator, but bosonic zero modes of the gauge fields. They are not removed by gauge transformations, and necessarily exist to cancel the anomaly inflow into each of the two intersecting 5-branes [1]. Each of small gauge rotation generators in $(\mathbf{27}, \mathbf{3}) \oplus (\overline{\mathbf{27}}, \overline{\mathbf{3}}) \oplus (\mathbf{1}, \mathbf{8})$ is an independent generator and gives rise to an independent zero mode. We also recall that exactly the same way of counting was done in the parallel symmetric 5-brane case, and was indeed consistent with the index analysis [10].

Next we examine the fermionic zero modes. The ten-dimensional heterotic gaugino equations of motion reads

$$\mathcal{D}(\omega - \frac{1}{3}H, A)\chi - \Gamma^M \chi \partial_M \phi + \frac{1}{8} \Gamma^M \gamma^{AB} (F_{AB} + \hat{F}_{AB})(\psi_M + \frac{2}{3} \Gamma_M \lambda) = 0, \quad (8)$$

where

$$D(\omega - \frac{1}{3}H, A)\chi \equiv \left(\partial_M + \frac{1}{4}(\omega_M^{AB} - \frac{1}{3}H_M^{AB})\Gamma_{AB} + \text{ad}A_M \right) \chi. \quad (9)$$

χ is in the adjoint **248** representation of E_8 , and $\text{ad}A_M \cdot \chi \equiv [A_M, \chi]$. If we set $\psi_M = 0$, $\lambda = 0$ and $\tilde{\chi} \equiv e^{-\phi} \chi$, then the equation is simplified to

$$\mathcal{D}(\omega - \frac{1}{3}H, A)\tilde{\chi} = 0. \quad (10)$$

Since there are no nontrivial backgrounds for the four-dimensional $i = 0, 7, 8, 9$ directions,

$$\Gamma^i \partial_i \tilde{\chi} + \Gamma^\mu D_\mu (\omega - \frac{1}{3}H, A)\tilde{\chi} = 0. \quad (11)$$

If $\tilde{\chi} = \tilde{\chi}_{4D} \otimes \tilde{\chi}_{6D}$, the second term is regarded as the mass term for the four-dimensional spinor $\tilde{\chi}_{4D}$. We are interested in the zero modes of this Dirac operator $\Gamma^\mu D_\mu (\omega - \frac{1}{3}H, A)$.

We fix the $SO(6)$ gamma matrices as

$$\begin{aligned} \gamma_1 &= \sigma_2 \otimes \mathbf{1} \otimes \mathbf{1}, & \gamma_2 &= \sigma_1 \otimes \sigma_1 \otimes \mathbf{1}, \\ \gamma_3 &= \sigma_1 \otimes \sigma_2 \otimes \mathbf{1}, & \gamma_4 &= \sigma_1 \otimes \sigma_3 \otimes \sigma_1, \\ \gamma_5 &= \sigma_1 \otimes \sigma_3 \otimes \sigma_2, & \gamma_6 &= \sigma_1 \otimes \sigma_3 \otimes \sigma_3. \end{aligned} \quad (12)$$

The six-dimensional chiral operator is $\gamma_\sharp \equiv -i\gamma_1\gamma_2\cdots\gamma_6 = \sigma_3 \otimes \mathbf{1} \otimes \mathbf{1}$. For $SO(9, 1)$ gamma matrices, we take $\Gamma^a = \gamma_{4D}^a \otimes \mathbf{1}_8$ ($a = 0, 7, 8, 9$), $\Gamma^\alpha = \gamma_{4D}^\sharp \otimes \gamma_\alpha$ ($\alpha = 1, \dots, 6$), where

γ_{4D}^a 's ($a = 0, 7, 8, 9$) are the ordinary $SO(3, 1)$ gamma matrices in the chiral representation. The ten-dimensional chirality is $\Gamma_{11} \equiv \gamma_{4D}^\# \otimes \gamma_\# = (\sigma_3 \otimes \mathbf{1}) \otimes (\sigma_3 \otimes \mathbf{1} \otimes \mathbf{1})$.

Now we consider the Dirac equation

$$\Gamma^\mu D_\mu(\omega - \frac{1}{3}H, A)\tilde{\chi} = 0. \quad (13)$$

The 16-component $SO(9, 1)$ (Majorana-)Weyl spinor χ (or $\tilde{\chi}$) is decomposed in terms of $SO(3, 1)$ and $SO(6)$ spinors as $\mathbf{16} = (\mathbf{2}_+, \mathbf{4}_+) \oplus (\mathbf{2}_-, \mathbf{4}_-)$, where the subscripts are the $SO(3, 1)$ and $SO(6)$ chiralities, $\gamma_{4D}^\#$ and $\gamma_\#$, respectively. Since $\tilde{\chi}$ is Majorana (but complex in this representation), the $(\mathbf{2}_+, \mathbf{4}_+)$ and $(\mathbf{2}_-, \mathbf{4}_-)$ components are not independent but are transformed each other by a charge conjugation.

As $\Gamma^\mu D_\mu(\omega - \frac{1}{3}H, A)$ is $SO(3, 1)$ diagonal, it is enough to consider

$$\gamma^\mu D_\mu(\omega - \frac{1}{3}H, A)\tilde{\chi}_{6D} = 0, \quad (14)$$

with the understanding that each component of $\tilde{\chi}_{6D}$ is accompanied by a two-component $SO(3, 1)$ Weyl spinor with a correlated chirality ($\gamma_\# \gamma_{4D}^\# = +1$).

On the other hand, we are interested in the gaugino zero modes in $(\mathbf{27}, \mathbf{3})$ or $(\overline{\mathbf{27}}, \overline{\mathbf{3}})$ in the decomposition $E_8 \supset E_6 \times SU(3)$ of $\mathbf{248}$. The gauge connections A_M take only nonzero values in the $SU(3)$ subalgebra, and we look for the zero modes $\tilde{\chi}_{6D}$ transforming as a triplet, either $\mathbf{3}$ or $\overline{\mathbf{3}}$, of $SU(3)$.

Since γ^α 's are in the form:

$$\begin{aligned} \gamma^1 &= \begin{pmatrix} & -i\mathbf{1} \otimes \mathbf{1} \\ -i\mathbf{1} \otimes \mathbf{1} & \end{pmatrix}, \\ \gamma^{\tilde{\alpha}} &= \begin{pmatrix} & \tilde{\gamma}^{\tilde{\alpha}} \\ \tilde{\gamma}^{\tilde{\alpha}} & \end{pmatrix} \quad (\tilde{\alpha} = 2, \dots, 6), \end{aligned} \quad (15)$$

and $\omega_\mu^{\alpha\beta}$, $H_\mu^{\alpha\beta}$ and $A_\mu^{\alpha\beta}$ all vanish if $\mu = 1$, (14) is reduced to two independent differential equations

$$\frac{i}{h} \frac{d}{dx^1} \tilde{\chi}_{6D}^+ + M^+ \tilde{\chi}_{6D}^+ = 0, \quad (16)$$

$$\frac{i}{h} \frac{d}{dx^1} \tilde{\chi}_{6D}^- - M^- \tilde{\chi}_{6D}^- = 0, \quad (17)$$

where $\tilde{\chi}_{6D}^\pm$ is the upper and lower components having definite chiralities: $\tilde{\chi}_{6D} = \begin{pmatrix} \tilde{\chi}_{6D}^+ \\ \tilde{\chi}_{6D}^- \end{pmatrix}$.

$\tilde{\chi}_{6D}^+$ ($\tilde{\chi}_{6D}^-$) is a **4** $SO(6)$ Weyl spinor, and each of the four components is a triplet of $SU(3)$.

Thus M^+ (M^-) is a $(4 \times 3 =) 12$ -by-12 matrix, given explicitly by

$$\begin{pmatrix} & M^- \\ M^+ & \end{pmatrix} \equiv \frac{h'}{h^2} \left(\begin{pmatrix} 0 & 0 & 0 & 0 & -\frac{3i}{2} & -\frac{i}{4} & 0 & -\frac{i}{4} \\ 0 & 0 & 0 & 0 & -\frac{i}{4} & -\frac{3i}{2} & -\frac{i}{4} & 0 \\ 0 & 0 & 0 & 0 & 0 & -\frac{i}{4} & -\frac{3i}{2} & -\frac{i}{4} \\ 0 & 0 & 0 & 0 & -\frac{i}{4} & 0 & -\frac{i}{4} & -\frac{3i}{2} \\ \frac{3i}{2} & -\frac{i}{4} & 0 & -\frac{i}{4} & 0 & 0 & 0 & 0 \\ -\frac{i}{4} & \frac{3i}{2} & -\frac{i}{4} & 0 & 0 & 0 & 0 & 0 \\ 0 & -\frac{i}{4} & \frac{3i}{2} & -\frac{i}{4} & 0 & 0 & 0 & 0 \\ -\frac{i}{4} & 0 & -\frac{i}{4} & \frac{3i}{2} & 0 & 0 & 0 & 0 \end{pmatrix} \otimes \mathbf{1}_3 \right. \\ \left. + \begin{pmatrix} 0 & 0 & 0 & 0 & -\frac{s\lambda_4}{2} & \frac{-s\lambda_1-\lambda_5}{2} & \frac{-2\lambda_2-s\lambda_9}{4} & 0 \\ 0 & 0 & 0 & 0 & \frac{\lambda_5-s\lambda_1}{2} & \frac{s\lambda_4}{2} & 0 & \frac{-2\lambda_2-s\lambda_9}{4} \\ 0 & 0 & 0 & 0 & \frac{2\lambda_2-s\lambda_9}{4} & 0 & \frac{s\lambda_4}{2} & \frac{s\lambda_1+\lambda_5}{2} \\ 0 & 0 & 0 & 0 & 0 & \frac{2\lambda_2-s\lambda_9}{4} & \frac{s\lambda_1-\lambda_5}{2} & -\frac{s\lambda_4}{2} \\ -\frac{s\lambda_4}{2} & \frac{-s\lambda_1-\lambda_5}{2} & \frac{-2\lambda_2-s\lambda_9}{4} & 0 & 0 & 0 & 0 & 0 \\ \frac{\lambda_5-s\lambda_1}{2} & \frac{s\lambda_4}{2} & 0 & \frac{-2\lambda_2-s\lambda_9}{4} & 0 & 0 & 0 & 0 \\ \frac{2\lambda_2-s\lambda_9}{4} & 0 & \frac{s\lambda_4}{2} & \frac{s\lambda_1+\lambda_5}{2} & 0 & 0 & 0 & 0 \\ 0 & \frac{2\lambda_2-s\lambda_9}{4} & \frac{s\lambda_1-\lambda_5}{2} & -\frac{s\lambda_4}{2} & 0 & 0 & 0 & 0 \end{pmatrix} \right), \quad (18)$$

where $\lambda_9 \equiv 3\lambda_3 + \sqrt{3}\lambda_8$. In identifying the spin connection as an $SU(3)$ gauge connection,

$s = \begin{pmatrix} & 1 \\ -1 & \end{pmatrix}$ can either be mapped to i , or to $-i$, and depending on this choice, the $SU(3)$

gauge connection matrix becomes one in the **3** representation, or in the $\bar{\mathbf{3}}$ representation.

Since $\tilde{\chi}_{6D}^+$ and $\tilde{\chi}_{6D}^-$ are not independent, we have only to solve the equation (16), and the solutions to (17) may then be obtained by a charge conjugation. To solve (16), we diagonalize M^+ to obtain its eigenvalues. Let $i\lambda$ be an eigenvalue of the *constant* matrix $(\frac{h'}{h^2})^{-1} M^+$, and $\psi_\lambda(x^1)$ be the corresponding eigenfunction, then they satisfy

$$\frac{i}{h} \psi'_\lambda + i\lambda \frac{h'}{h^2} \psi_\lambda = 0. \quad (19)$$

This is solved to give

$$\psi_\lambda(x^1) = \text{const.} (h(x^1))^{-\lambda}. \quad (20)$$

Thus, for each eigenvalue, there exists a zero mode of the Dirac operator. Since ξ is negative for positive tension, if $\lambda < 1$, the mode is localized near $x^1 = 0$, while if $\lambda \geq 1$, it is not localized, being either non-normalizable or localized rather at “infinity” $x^1 = \pm \frac{h_0}{|\xi|}$.

The list of eigenvalues of $(\frac{h'}{h^2})^{-1} M^+$ is as follows: If $s = +i$, the eigenvalues are

$$\left\{ 2i, \frac{3i}{2}, \frac{3i}{2}, i, -i, i, i, i, \frac{3i}{2}, \frac{3i}{2}, \frac{7i}{2}, \frac{7i}{2} \right\}, \quad (21)$$

while if $s = -i$, they are

$$\left\{ 2i, \frac{3i}{2}, \frac{3i}{2}, i, 2i, 4i, 2i, 2i, -\frac{i}{2}, -\frac{i}{2}, \frac{3i}{2}, \frac{3i}{2} \right\}. \quad (22)$$

We can clearly see an asymmetry between (21) and (22), in particular that the former has only one negative (times imaginary unit) eigenvalue, while the latter has two negative eigenvalues. Assuming that the branes have positive tension so that the function $h(x)$ has the profile shown in FIG.2, these are the only modes whose profiles have a peak at $x^1 = 0$ or $z = 0$ in the coordinate (4). The same is also true for the original gaugino variable $\chi = h\tilde{\chi}$ (although the modes with $\lambda = +1$ then become constant). This result implies that there are indeed three localized modes, and two of them are in one (say, $(\mathbf{27}, \mathbf{3})$) representation, and the rest belongs to the other $((\overline{\mathbf{27}}, \overline{\mathbf{3}}))$ representation.

Acknowledgments

We would like to thank Tohru Eguchi, Satoshi Iso, Hikaru Kawai, Taichiro Kugo, Hiroshi Kunitomo, Nobuyoshi Ohta and Shigeki Sugimoto for illuminating discussions. We are also grateful to Keiichi Akama, Masafumi Fukuma, Machiko Hatsuda, Takeo Inami, Akihiro Ishibashi, Katsushi Ito, Katsumi Itoh, Hiroshi Itoyama, Yoichi Kazama, Yoshio Kikukawa, Yoshihisa Kitazawa, Hideo Kodama, Nobuhiro Maekawa, Nobuhito Maru, Yoji Michishita, Muneto Nitta, Kazutoshi Ohta, Soo-Jong Rey, Tomohiko Takahashi, Shinya Tomizawa, Tamiaki Yoneya and Kentaro Yoshida for discussions and comments. We thank YITP for hospitality and support during the workshop: “Branes, Strings and Black Hole”, where part of this work was done. This work is supported by Grant-in-Aid for Scientific Research (C) #20540287-H20 from The Ministry of Education, Culture, Sports, Science and Technology of Japan.

[1] T. Kimura and S. Mizoguchi, “*Chiral Generations on Intersecting 5-branes in Heterotic String Theory*,” arXiv:0912.1334 [hep-th], KEK-TH-1341.

- [2] R. Argurio, F. Englert and L. Houart, Phys. Lett. B **398**, 61 (1997) [arXiv:hep-th/9701042]
N. Ohta, Phys. Lett. B **403**, 218 (1997) [arXiv:hep-th/9702164].
- [3] S. J. Rey, Phys. Rev. D **43**, 526 (1991); “*Axionic String Instantons And Their Low-Energy Implications*,” Invited talk given at Workshop on Superstrings and Particle Theory, Tuscaloosa, Alabama, Nov 8-11, 1989. Published in Tuscaloosa Workshop 1989:0291-300; “*On string theory and axionic strings and instantons*,” Presented at Particle and Fields ’91 Conf., Vancouver, Canada, Aug 18-22, 1991. Published in DPF Conf.1991:0876-881.
- [4] A. Strominger, Nucl. Phys. B **343**, 167 (1990) [Erratum-ibid. B **353**, 565 (1991)].
- [5] C. G. Callan, J. A. Harvey and A. Strominger, Nucl. Phys. B **359**, 611 (1991); Nucl. Phys. B **367**, 60 (1991).
- [6] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999) [arXiv:hep-ph/9905221]; Phys. Rev. Lett. **83**, 4690 (1999) [arXiv:hep-th/9906064].
- [7] E. A. Bergshoeff and M. de Roo, Nucl. Phys. B **328**, 439 (1989);
- [8] T. Kimura and P. Yi, JHEP **0607**, 030 (2006) [arXiv:hep-th/0605247].
- [9] T. Kimura, JHEP **0708**, 048 (2007) [arXiv:0704.2111 [hep-th]].
- [10] D. Bellisai, Nucl. Phys. B **467**, 127 (1996) [arXiv:hep-th/9511198].
- [11] This corrects the statement made in the earlier version of this paper, in which it was erroneously conjectured that the three supermultiplets would be of the same chirality.
- [12] More curiously, although the branes have a positive tension as we have derived (3), the 4D metric is inversely warped (like near the negative tension brane in the RS I model) in the Einstein frame.